

Aanvullende informatie bij Bijenhouden februari 2014

Referenties bij '**Jaarrond bestrijden Varroamijt vermindert wintersterfte**', J. van der Steen e.a., Bijenhouden februari 2014 pag. 12 en 13.

- Elke Genersch et al., 2010, The German bee monitoring project: a long term study to understand periodically high winter losses of honey bee colonies* Apidologie 41: 332–352, DOI: [10.1051/apido/2010014](https://doi.org/10.1051/apido/2010014)
- Ernesto Guzman-Novoa, et al., 2010. *Varroa destructor* is the main culprit for the death and reduced populations of overwintered honey bee (*Apis mellifera*) colonies in Ontario, Canada. Apidologie 41: 443–450, DOI: [10.1051/apido/2009076](https://doi.org/10.1051/apido/2009076)
- Marc O. Schäfer, et al., 2010, Winter Losses of Honeybee Colonies (Hymenoptera: Apidae): The Role of Infestations with *Aethina tumida* (Coleoptera: Nitidulidae) and *Varroa destructor* (Parasitiformes: Varroidae). J. Econ. Entomol. 103(1):10–16 (2010); DOI: 10.1603/EC09233
- Yves Le Conte et al., 2010. *Varroa* mites and honey bee health: can *Varroa* explain part of the colony losses? Apidologie 41:353–363, DOI: [10.1051/apido/2010017](https://doi.org/10.1051/apido/2010017)
- Vanengelsdorp, D. et al., 2013. Standard epidemiological methods to understand and improve *Apis mellifera* health. In V Dietemann; J D Ellis, P Neumann (Eds) The COLOSS BEEBOOK: Volume II: Standard methods for *Apis mellifera* pest and pathogen research. Journal of Apicultural Research 52(1): <http://dx.doi.org/10.3896/IBRA.1.52.1.08>

Literatuur bij '**Ook een honingbij ontgift zich**', Henk van der Scheer en Tjeerd Blacquièrè, Bijenhouden februari 2014, pag. 29-31

- Belzunces, L.P. e.a., 2012. Neural effects of insecticides in the honey bee. Apidologie 43:348–370.
- Blacquièrè, T. e.a., 2012. Neonicotinoids in bees: a review on concentrations, side-effects and risk assessment. Ecotoxicology 21:973-992.
- Casida, J.E. en Durkin, K.A., 2013. Neuroactive insecticides: targets, selectivity, resistance, and secondary effects. Annual Review of Entomology 58:99-117.
- Claudianos, C. e.a., 2006. A deficit of detoxification enzymes: pesticide sensitivity and environmental response in the honeybee. Insect Molecular Biology 15(5):615-636.
- Cresswell, J.E. e.a., 2012. Differential sensitivity of honey bees and bumble bees to a dietary insecticide (imidacloprid). Zoology 115(6):365-371.
- Cresswell, J.E. e.a., 2013. Clearance of ingested neonicotinoid pesticide (imidacloprid) in honey bees (*Apis mellifera*) and bumble bees (*Bombus terrestris*). Pest Management Science.. doi:10.1002/ps.3569
- Decourtye, A. en Devillers, J., 2010. Ecotoxicity of neonicotinoid insecticides to bees. Hoofdstuk 8 in 'Insect nicotinic acetylcholine receptors', vol. 683, pp. 85-95, uit de serie: Advances in Experimental Medicine and Biology.
- Derecka, K. e.a., 2013. Transient exposure to low levels of insecticide affects metabolic networks of honeybee larvae. PLoS ONE 8(7): e68191.
- Hawthorne, D.J. en Dively, G.P., 2011. Killing them with kindness? In-hive medications may inhibit xenobiotic efflux transporters and endanger honey bees PLoS ONE 6(11): e26796.

- Honeybee Genome Sequencing Consortium (Overall project leadership: George M. Weinstock en Gene E. Robinson), 2006. Insights into social insects from the genome of the honeybee *Apis mellifera*. *Nature* 443:931-949.
- Mao, W. e.a., 2011. CYP9Q-mediated detoxification of acaricides in the honey bee (*Apis mellifera*). *Proceedings of the National Academy of Sciences USA* 108(31):12657-12662.
- Medrzycki, P. e.a., 2003. Effects of imidacloprid administered in sub-lethal doses on honey bee behaviour. Laboratory tests. *Bulletin of Insectology* 56(1):59-62.
- Mullin, C.A. e.a., 2010. High levels of miticides and agrochemicals in North American apiaries: implications for honey bee health. *Plos One* 5(3):e9754.
- Schneider, C.W. e.a., 2012. RFID Tracking of sublethal effects of two neonicotinoid insecticides on the foraging behavior of *Apis mellifera*. *PLoS ONE* 7(1): e30023.
- Suchail, S. e.a., 2003. Metabolism of imidacloprid in *Apis mellifera*. *Pest Management Science* 60(3):291-296.
- Suchail, S. e.a., 2004. In vivo distribution and metabolisation of 14C-imidacloprid in different compartments of *Apis mellifera* L. *Pest Management Science* 60(11):1056-1062.
- Tomizawa, M. en Casida, J.E., 2003. Selective toxicity of neonicotinoids attributable to specificity of insect and mammalian nicotinic receptors. *Annual Review of Entomology* 48: 339-364.
- Wallner, K., 1999. Bienenschäden im Raps? *ADIZ* 33(5):7-10.
- Yang, E-C. e. a., 2012. Impaired olfactory associative behavior of honeybee workers due to contamination of imidacloprid in the larval stage. *PLoS ONE* 7(11): e49472]